



**LGV CAN CAUSE
MAJOR INFLAMMATION
INSIDE THE ASS.**

**MOST AT RISK ARE
MEN FUCKING WITHOUT
CONDOMS OR FISTING
WITHOUT LATEX GLOVES.**

**EASILY TREATED
WITH ANTIBIOTICS.**

**LEFT UNTREATED CAN END
IN SERIOUS DAMAGE TO
THE ASS AND GENITALS.**

ACT

AIDS Committee of Toronto

399 Church Street, 4th floor

Toronto, ON M5B 2J6

Tel: 416-340-2437

Email: ask@actontario.org

Website: www.actontario.org/lgv

Adapted and reprinted with permission
of Terrence Higgins Trust

Design Felton Communication +44(0)20 7405 0900

Ref: 9742 Photos courtesy of Millivres-Prowler Group



LGV

AN INFECTION GROWING AMONG GAY MEN

WHAT IS LGV?

LGV stands for 'lymphogranuloma venereum'. It's a sexually transmitted infection caused by a certain type of chlamydia bacteria. It's started to spread among gay men so we need to be looking out for it.

IS THIS NEW?

It's common in places like Africa, Asia and the Caribbean. There have been outbreaks among gay men in some European cities; some gay men in the United States and Canada have also been recently diagnosed with LGV.

WHO GETS LGV?

So far the infection's been seen mostly in gay men who:

- **FUCK OR GET FUCKED WITHOUT CONDOMS**
- **FIST WITHOUT GLOVES**
- **SUCK OR GET SUCKED WITHOUT CONDOMS**

Most of the men infected so far have been HIV positive.

HOW IS LGV PASSED ON?

The bacteria get into someone's body through the warm moist skin of the mouth, cock or inside of the ass. **Fucking without condoms** and **fisting without gloves** are the easiest way this happens. Using things like dildos on more than one man's ass could spread LGV too.

WHAT HAPPENS IF I GET LGV?

Most gay men getting LGV recently had the infection inside their ass, causing major inflammation there. But LGV has three stages.

STAGE 1: Three days to three weeks after getting infected there may be a small, painless sore where the bacteria got into your mouth, cock or inside your ass. Many people don't get a sore or don't notice it. You will be able to give LGV to others from this point until you get treated.

STAGE 2: 10-30 days later (or even longer) your glands may become painfully swollen. You might feel ill and have a fever.

LGV in the ass - You can get painful inflammation in your ass (called 'proctitis') and blood or pus coming from it. Shitting may be painful, you might get constipated or develop painful abscesses/ulcers in the ass.

LGV in the cock - You might get a discharge, the glands in your groin can swell - so much that they might push through your skin and leak fluid.

LGV in the mouth - If infected through the mouth or throat, glands in your neck or armpit may swell.

STAGE 3: Left untreated LGV can later go on to cause serious problems. Haemorrhoid-like growths can appear, your cock and balls can swell massively and the inside of your ass can be seriously damaged; you may need to have surgery.

CAN LGV BE CURED?

As long as it's treated before it gets to the third stage, LGV is quickly cured with 3 weeks of antibiotics and leaves no lasting damage.

HOW CAN I PROTECT MYSELF?

There's no vaccine and if you've had LGV once you can get it again. Condoms reduce the chances of getting LGV, as do latex gloves during fisting. If several men are being fucked or fisted in the same session, condoms and gloves should be changed with each man (to stop the infection being spread from man to man). Dildos etc should not be used on more than one ass in one sex session - or should be covered with a condom that can be changed with each ass it's used on.

If you have LGV, samples will test positive for chlamydia but more tests will be needed to be sure if you have LGV and not the more common type of chlamydia. A check-up is crucial if you have any symptoms of inflammation inside your ass - or in your cock, mouth or throat. Sex should be avoided until you've been checked.

WHAT ELSE DO I NEED TO KNOW?

Having LGV makes it easier for you to become infected by HIV and to pass on HIV to your partner because of the bleeding and skin damage LGV causes. It also puts you at increased risk for other STIs (such as syphilis), and other infections like Hepatitis B and C which can cause serious liver disease.